

§ 180.436**40 CFR Ch. I (7-1-10 Edition)**

(i) In or on all food/feed items (other than those covered by a higher tolerance as a result of use on growing crops) in food/feed handling establishments.

(ii) The insecticide may be present as a residue from application of deltamethrin in food handling establishments, including food service, manufacturing and processing establishments, such as restaurants, cafeterias, supermarkets, bakeries, breweries, dairies, meat slaughtering and packing plants, and canneries, feed handling establishments including feed manufacturing and processing establishments, in accordance with the following prescribed conditions:

(A) Application shall be limited to general surface and spot and/or crack and crevice treatment in food/feed handling establishments where food/feed and food/feed products are held, processed, prepared and served. General surface application may be used only when the facility is not in operation provided exposed food/feed has been covered or removed from the area being treated. Spot and/or crack and crevice application may be used while the facility is in operation provided exposed food/feed is covered or removed from the area being treated prior to application. Spray concentration shall be limited to a maximum of 0.06 percent active ingredient. Contamination of food/feed or food/feed contact surfaces shall be avoided.

(B) To assure safe use of the insecticide, its label and labeling shall conform to that registered with the U.S. Environmental Protection Agency and shall be used in accordance with such label and labeling.

(b) *Section 18 emergency exemptions.* [Reserved]

(c) *Tolerances with regional registrations.* [Reserved]

(d) *Indirect or inadvertent residues.* [Reserved]

[62 FR 63001, Nov. 26, 1997, as amended at 63 FR 45414, Aug. 26, 1998; 69 FR 62614, Oct. 27, 2004; 74 FR 46375, Sept. 9, 2009]

§ 180.436 Cyfluthrin and the isomer beta-cyfluthrin; tolerances for residues.

(a) *General.* (1) Tolerances are established for residues of the insecticide

cyfluthrin (cyano(4-fluoro-3-phenoxyphenyl)methyl-3-(2,2-dichloroethenyl)-2,2dimethylcyclopropane-carboxylate; CAS No. 68359-37-5) in or on the following raw agricultural commodities:

Commodity	Parts per million
Alfalfa	5.0
Alfalfa, forage	5.0
Alfalfa, hay	13
Almond, hulls	0.5
Barley, bran	0.5
Barley, grain	0.15
Beet, sugar, dried pulp	1.0
Beet, sugar, roots	0.10
Brassica, head and stem, subgroup 5A	2.5
Brassica, leafy greens, subgroup 5B	7.0
Buckwheat, grain	0.15
Carrot, roots	0.20
Cattle, fat	2.0
Cattle, meat	0.10
Cattle, meat byproducts	0.10
Citrus, dried pulp	0.3
Citrus, oil	0.3
Corn, field, grain	0.05
Corn, pop, grain	0.05
Corn, sweet, kernel plus cob with husks removed	0.05
Cotton, hulls	2.0
Cotton, refined oil	2.0
Cotton, undelinted seed	1.0
Egg	0.01
Fruit, citrus, group 10	0.2
Fruit, pome, group 11	0.5
Fruit, stone, group 12	0.3
Goat, fat	2.0
Goat, meat	0.05
Goat, meat byproducts	0.05
Grain, aspirated fractions	150
Grain, cereal, forage, fodder and hay, group 16, forage, except rice	25
Grain, cereal, forage, fodder and hay, group 16, hay, except rice	6.0
Grain, cereal, forage, fodder and hay, group 16, stover, except rice	30
Grain, cereal, forage, fodder and hay, group 16, straw, except rice	7.0
Grape	1.0
Grape, raisin	3.5
Grass, forage, fodder and hay, group 17, forage	12
Grass, forage, fodder and hay, group 17, hay	50
Hog, fat	0.5
Hog, meat	0.01
Hog, meat byproducts	0.01
Hop, dried cones	20.0
Hop, vines	4.0
Horse, fat	2.0
Horse, meat	0.05
Horse, meat byproducts	0.05
Lettuce, head	2.0
Lettuce, leaf	3.0
Milk	0.2
Milk, fat	5.0
Millet, grain	0.15
Mustard greens	7.0
Nut, tree, group 14	0.01
Oat, bran	0.5
Oat, grain	0.15
Pea, and bean, dried shelled, except soybean, subgroup 6C	0.15
Pea, dry, seed	0.15
Pea, southern, succulent	0.25

Environmental Protection Agency
§ 180.436

Commodity	Parts per million
Peanut	0.01
Peanut, hay	6.0
Pepper	0.50
Pistachio	0.01
Poultry, fat	0.01
Poultry, meat	0.01
Poultry, meat byproducts	0.01
Radish, roots	1.0
Rye, bran	0.5
Rye, grain	0.15
Sheep, fat	2.0
Sheep, meat	0.05
Sheep, meat byproducts	0.05
Sorghum, grain, grain	3.5
Soybean, forage	8.0
Soybean, hay	4.0
Soybean, seed	0.03
Sugarcane, cane	0.05
Sugarcane, molasses	0.20
Sunflower, forage	5.0
Sunflower, seed	0.02
Teosinte, grain	0.05
Tomato	0.20
Tomato, dry pomace	5.0
Tomato, paste	0.5
Tomato, wet pomace	5.0
Triticale, grain	0.15
Turnip, greens	7.0
Vegetable, cucurbit, group 9	0.1
Vegetable, fruiting, group 8	0.5
Vegetable, leafy, except brassica, group 4	6.0
Vegetable, tuberous and corm, subgroup 1C	0.01
Wheat, bran	0.5
Wheat, grain	0.15
Wheat, shorts	0.5

(2) A tolerance of 0.05 ppm is established for residues of the insecticide cyfluthrin (cyano(4-fluoro-3-phenoxyphenyl)methyl-3-(2,2-dichloroethenyl)-2,2-dimethylcyclopropanecarboxylate; CAS Reg. No. 69359-37-5) in food commodities exposed to the insecticide during treatment of food-handling establishments where food and food products are held, processed, prepared, or served. Treatments may be made by general surface, spot, and/or crack and crevice applications.

(i) General surface treatments shall be limited to a maximum of 3.8 grams of active ingredient per 1,000 square feet, applying to walls, floors, and ceilings with a low-pressure system. Cover or remove all food processing and/or handling equipment during application. Do not apply directly to food products. Reapplications may be made at 10-day intervals.

(ii) Crack and crevice or spot treatments shall be limited to a maximum of 0.1 percent of the active ingredient weight, applied with a low-pressure system with a pinpoint or variable-pat-

tern nozzle. Dust formulation shall be limited to a maximum of 0.1 percent of the active ingredient by weight, applied using a hand duster, power duster, or other equipment capable of applying dust insecticide directly into voids and cracks and crevices. Dust applications should be made in a manner to avoid deposits on exposed surfaces or introducing the material into the air. Cover exposed food or remove food from premises. Do not apply directly to food. Reapplications may be made at 10-day intervals.

(iii) To ensure safe use of the insecticide, its label and labeling shall conform to that registered by the Environmental Protection Agency, and it shall be used in accordance with such label and labeling.

(3) A tolerance of 0.05 part per million is established for residues of the insecticide cyfluthrin (cyano(4-fluoro-3-phenoxyphenyl)methyl-3-(2,2-dichloroethenyl)-2,2-dimethylcyclopropanecarboxylate; CAS Reg. No. 69359-37-5) in feed commodities exposed to the insecticide during treatment of feed-handling establishments where feed and feed products are held, processed, prepared, or served. Treatments may be made by general surface, spot, and/or crack and crevice applications.

(i) General surface treatments shall be limited to a maximum of 3.8 grams of active ingredient per 1,000 square feet, applying to walls, floors, and ceilings with a low-pressure system. Cover or remove all feed processing and/or handling equipment during application. Do not apply directly to feed products. Reapplications may be made at 10-day intervals.

(ii) Crack and crevice or spot treatments shall be limited to a maximum of 0.1 percent of the active ingredient by weight, applied with a low-pressure system with a pinpoint or variable-pattern nozzle. Dust formulation shall be limited to a maximum of 0.1 percent of the active ingredient by weight, applied using a hand duster, power duster, or other equipment capable of applying dust insecticide directly into voids and cracks and crevices. Dust applications should be made in a manner to avoid deposits on exposed surfaces or introducing the material into the

§ 180.436

air. Cover exposed feed or remove feed from premises. Do not apply directly to feed. Reapplications may be made at 10-day intervals.

(iii) To ensure safe use of the insecticide, its label and labeling shall conform to that registered by EPA, and it shall be used in accordance with such label and labeling.

(4) Tolerances are established for residues of the isomer, beta-cyfluthrin, cyano(4-fluoro-3-phenoxyphenyl)methyl-3-(2,2-dichloroethenyl)-2,2-dimethylcyclopropanecarboxylate [mixture comprising the enantiomeric pair (*R*)- α -cyano-4-fluoro-3-phenoxybenzyl (1*S*,3*S*)-3-(2,2-dichlorovinyl)-2,2-dimethylcyclopropanecarboxylate and (*S*)- α -cyano-4-fluoro-3-phenoxybenzyl (1*R*,3*R*)-3-(2,2-dichlorovinyl)-2,2-dimethylcyclopropanecarboxylate with the enantiomeric pair (*R*)- α -cyano-4-fluoro-3-phenoxybenzyl (1*S*,3*R*)-3-(2,2-dichlorovinyl)-2,2-dimethylcyclopropanecarboxylate and (*S*)- α -cyano-4-fluoro-3-phenoxybenzyl (1*R*,3*S*)-3-(2,2-dichlorovinyl)-2,2-dimethylcyclopropanecarboxylate], in or on the following raw agricultural commodities:

Commodity	Parts per million
Alfalfa	5.0
Alfalfa, forage	5.0
Alfalfa, hay	13
Almond, hulls	0.5
Barley, bran	0.5
Barley, grain	0.15
Beet, sugar, dried pulp	1.0
Beet, sugar, roots	0.10
Brassica, head and stem, subgroup 5A	2.5
Brassica, leafy greens, subgroup 5B	7.0
Buckwheat, grain	0.15
Carrot, roots	0.20
Cattle, fat	2.0
Cattle, meat	0.10
Cattle, meat byproducts	0.10
Citrus, dried pulp	0.3
Citrus, oil	0.3
Corn, field, grain	0.05
Corn, pop, grain	0.05
Corn, sweet, kernel plus cob with husks removed	0.05
Cotton, hulls	4
Cotton, refined oil	2.0
Cotton, undelinted seed	2.0
Egg	1.0
Fruit, citrus, group 10	0.01
Fruit, pome, group 11	0.2
Fruit, stone, group 12	0.5
Goat, fat	0.3
Goat, meat	2.0
Goat, meat byproducts	0.05
Grain, aspirated fractions	0.05

150

40 CFR Ch. I (7-1-10 Edition)

Commodity	Parts per million
Grain, cereal, forage, fodder and hay, group 16, forage, except rice	25
Grain, cereal, forage, fodder and hay, group 16, hay, except rice	6.0
Grain, cereal, forage, fodder and hay, group 16, stover, except rice	30
Grain, cereal, forage, fodder and hay, group 16, straw, except rice	7.0
Grape	1.0
Grape, raisin	3.5
Grass, forage, fodder and hay, group 17, forage	12
Grass, forage, fodder and hay, group 17, hay	50
Hog, fat	0.5
Hog, meat	0.01
Hog, meat byproducts	0.01
Hop, dried cones	20.0
Hop, vines	4.0
Horse, fat	2.0
Horse, meat	0.05
Horse, meat byproducts	0.05
Lettuce, head	2.0
Lettuce, leaf	3.0
Milk	0.2
Milk, fat	5.0
Millet, grain	0.15
Mustard greens	7.0
Nut, tree, group 14	0.01
Oat, bran	0.5
Oat, grain	0.15
Pea and bean, dried shelled, except soybean, subgroup 6C	0.15
Pea, dry, seed	0.15
Pea, southern, succulent	0.25
Peanut	0.01
Peanut, hay	6.0
Pepper	0.50
Pistachio	0.01
Poultry, fat	0.01
Poultry, meat	0.01
Poultry, meat byproducts	0.01
Radish, roots	1.0
Rye, bran	0.5
Rye, grain	0.15
Sheep, fat	2.0
Sheep, meat	0.05
Sheep, meat byproducts	0.05
Sorghum, grain, grain	3.5
Soybean, forage	8.0
Soybean, hay	4.0
Soybean, seed	0.03
Sugarcane, cane	0.05
Sugarcane, molasses	0.20
Sunflower, forage	5.0
Sunflower, seed	0.02
Teosinte, grain	0.05
Tomato	0.20
Tomato, paste	0.5
Tomato, pomace	5.0
Triticale, grain	0.15
Turnip, greens	7.0
Vegetable, cucurbit, group 9	0.1
Vegetable, fruiting, group 8	0.5
Vegetable, leafy greens, except Brassica, group 4	6.0
Vegetable, tuberous and corm, subgroup 1C	0.01
Wheat, bran	0.5
Wheat, grain	0.15
Wheat, shorts	0.5

(b) Section 18 emergency exemptions.

[Reserved]

(c) Tolerances with regional registrations. [Reserved]

Environmental Protection Agency
§ 180.438

(d) *Indirect or inadvertent residues.*
 [Reserved]

[53 FR 1924, Jan. 25, 1988]

EDITORIAL NOTE: For FEDERAL REGISTER citations affecting § 180.436, see the List of CFR Sections Affected, which appears in the Finding Aids section of the printed volume and on GPO Access.

§ 180.437 Methyl 2-(4-isopropyl-4-methyl-5-oxo-2-imidazolin-2-yl)-*p*-toluate and methyl 6-(4-isopropyl-4-methyl-5-oxo-2-imidazolin-2-yl)-*m*-toluate; tolerances for residues.

Tolerances are established for the combined residues of the herbicide methyl 2-(4-isopropyl-4-methyl-5-oxo-2-imidazolin-2-yl)-*p*-toluate and methyl 6-(4-isopropyl-4-methyl-5-oxo-2-imidazolin-2-yl)-*m*-toluate in or on the following raw agricultural commodities:

Commodity	Parts per million
Barley, grain	0.10
Barley, straw	2.00
Sunflower, seed	0.10
Wheat, grain	0.10
Wheat, straw	2.00

[53 FR 24069, June 27, 1988]

§ 180.438 Lambda-cyhalothrin and an isomer gamma-cyhalothrin; tolerances for residues.

(a) *General.* (1) Tolerances are established for the combined residues of the pyrethroid lambda-cyhalothrin, 1:1 mixture of (*S*)- α -cyano-3-phenoxybenzyl-(*Z*)-(1*R*,3*R*)-3-(2-chloro-3,3,3-trifluoroprop-1-enyl)-2,2-dimethylcyclopropanecarboxylate and (*R*)- α -cyano-3-phenoxybenzyl-(*Z*)-(1*S*,3*S*)-3-(2-chloro-3,3,3-trifluoroprop-1-enyl)-2,2-dimethylcyclopropanecarboxylate and its epimer expressed as epimer of lambda-cyhalothrin, a 1:1 mixture of (*S*)- α -cyano-3-phenoxybenzyl-(*Z*)-(1*S*,3*S*)-3-(2-chloro-3,3,3-trifluoroprop-1-enyl)-2,2-dimethylcyclopropanecarboxylate and (*R*)- α -cyano-3-phenoxybenzyl-(*Z*)-(1*R*,3*R*)-3-(2-chloro-3,3,3-trifluoroprop-1-enyl)-2,2-dimethylcyclopropanecarboxylate, on plants and livestocks, as indicated in the following table.

Commodity	Parts per million
Alfalfa, forage	5.0
Alfalfa, hay	6.0
Almond, hulls	1.5
Apple, wet pomace	2.50
Avocado, imported	0.20
Barley, bran	0.2
Barley, grain	0.05
Barley, hay	2.0
Barley, straw	2.0
Brassica, head and stem, subgroup 5A	0.4
Buckwheat, grain	0.05
Canola, refined oil	2.0
Canola, seed	1.0
Cattle, fat	3.0
Cattle, meat	0.2
Cattle, meat byproducts	0.2
Corn, field, flour	0.15
Corn, field, forage	6.0
Corn, field, grain	0.05
Corn, field, stover	1.0
Corn, pop, grain	0.05
Corn, pop, grain, flour	0.05
Corn, pop, stover	1.0
Corn, sweet, forage	6.0
Corn, sweet, stover	1.0
Corn, sweet, kernel plus cob with husks removed	0.05
Cotton, undelinted seed	0.05
Egg	0.01
Fruit, pome, group 11	0.30
Fruit, stone, group 12	0.50
Garlic	0.1
Goat, fat	3.0
Goat, meat	0.2
Goat, meat byproducts	0.2
Grain, aspirated fractions	2.0
Grass, forage, fodder and hay, group 17	7.0
Hog, fat	0.2
Hog, meat	0.01
Hog, meat byproducts	0.02
Hop, dried cones	10.0
Horse, fat	3.0
Horse, meat	0.2
Horse, meat byproducts	0.2
Lettuce, head	2.0
Lettuce, leaf	2.0
Milk, fat (reflecting 0.4 ppm in whole milk)	10.0
Nut, tree, group 14	0.05
Oat, grain	0.05
Oat, forage	2.0
Oat, hay	2.0
Oat, straw	2.0
Onion, bulb	0.1
Pea and bean, dried shelled, except soybean, subgroup 6C	0.10
Pea and bean, succulent shelled, subgroup 6B	0.01
Peanut	0.05
Peanut, hay	3.0
Pistachio	0.05
Poultry, fat	0.03
Poultry, meat	0.01
Poultry, meat byproducts	0.01
Rice, grain	1.0
Rice, hulls	5.0
Rice, straw	1.8
Rice, wild, grain	1.0
Rye, bran	0.2
Rye, grain	0.05
Rye, forage	2.0
Rye, straw	2.0
Sheep, fat	3.0
Sheep, meat	0.2
Sheep, meat byproducts	0.2